

## GENERAL INFORMATION

KEYLATE Cobalt contains chelated complexes of environmentally active cobalt. Apply KEYLATE Cobalt to the soil or crop foliage to increase microorganism activity within the crop rhizosphere. KEYLATE Cobalt promotes robust populations of soil flora. Apply KEYLATE Cobalt before planting or early in plant development to maximize microbial life and promote plant vigor.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product inconsistent with its labeling.

KEYLATE Cobalt can be applied in-furrow, as a side-dress or foliar application, and through flood, drip, overhead, pivot and micro irrigation systems. Schedule regular applications of KEYLATE Cobalt or apply during crop transitional stages when root stress is most likely to occur. The addition of ammonium sulfate or equivalent at 0.5% v/v will increase the activity of KEYLATE Cobalt. Apply the recommended amount in a volume sufficient to provide uniform coverage.

**RECOMMENDED USES:** Apply to the soil or foliage starting with the transplant water or within 7 days of transplant or when seeded crops have 2 to 4 true leaves and continue through harvest.

For SOIL APPLICATION apply KEYLATE Cobalt at 1 to 4 pints per acre.

For FOLIAR APPLICATIONS apply KEYLATE Cobalt at 1 pint per acre on actively growing crops in a minimum of 20 GPA using split-applications when appropriate.

Use KEYLATE Cobalt on the following crops:

Apply every 7 to 21 days on Field Crops: alfalfa, bean crops, canola, clover, corn, cotton, forage grasses, peanut, potato, rape (canola), rice, sorghum, soybean, sugar beet, sugarcane, sunflower, tobacco, and wheat/barley/oat. Vegetable Crops: artichoke, asparagus, beet, carrot, cruciferous, fennel, herbs, leafy vegetables, melon, onion/garlic, cucurbits, eggplant, pepper, strawberry, sweet potato and tomato. Field Grown Nursery Stock: caneberries, strawberry, tree and vine crops and other grafted, clonal or vegetatively propagated stock plants.

Apply KEYLATE Cobalt every 7 to 21 days until 10% bloom or after petal fall to Fruit and Nut Crops: almond, apple, apricot, avocado, berry crops, cherry, citrus, date palm, fig, filbert, grapes, hazelnut, hops, nectarines, oil palm, olive, persimmon, peach, pear, pecan, pistachio, plums/prune, pomegranate, subtropical fruit and walnut.

FOR TRANSPLANTS AND HIGH VOLUME DRENCH:

Apply 4 pints of KEYLATE Cobalt per 500 gallons of water.

FOR AERIAL AND LOW VOLUME APPLICATION EQUIPMENT:

Apply 1 to 4 pints of KEYLATE Cobalt in a minimum of 5 GPA. Do not exceed 2 pints of KEYLATE Cobalt per gallon of total solution volume.

**COMPATIBILITY:** KEYLATE Cobalt is tank-mix compatible with most pesticides and fertilizers. Add KEYLATE Cobalt to the diluted solution last.

Always conduct a compatibility test for new or untried tank-mix combinations.

Conditions Of Sale: 1. Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an officer of Seller is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

## KEYLATE® Cobalt

### Guaranteed Analysis

0-0-0

**Cobalt (Co)** 5%  
5% Chelated Cobalt

Derived from cobalt sulfate, chelated with ethanol,  
2-amino-2-hydroxy-1,2,3-propanetricarboxylate

### CAUTION

**Keep out of Reach of Children  
May Cause Eye Irritation**

### USER SAFETY RECOMMENDATIONS

Avoid contact with skin, eyes or clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove wet clothing immediately. Wash skin thoroughly and put on clean clothing.

### FIRST AID

If in eyes: Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. If on skin: Take off contaminated clothing and rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. Have product container or label with you when calling a poison control center, or doctor, or going for treatment.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

### STORAGE AND DISPOSAL

Store this product out of direct sunlight, in a cool place above 40°F. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Dispose of empty container in a sanitary landfill or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals>.

NET WEIGHT: 10.35 lbs/gal (1.24 Kg/L) at 68°F

### NET CONTENTS:

- |  |   |
|--|---|
| <input type="checkbox"/> 2.5 gallons/25.9 lbs (11.75 Kg) | <input type="checkbox"/> 10 liter/12.4 kg |
| <input type="checkbox"/> 5.0 gallons/51.8 lbs (23.5 Kg)  | <input type="checkbox"/> 20 liter/24.8 kg |

# 5% COBALT



**Z-KEYCO**

**US Patent: 5,997,600**



**StollerUSA**

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway #400 | Houston Texas 77055 USA

Phone: 1-800-539-5283 or 713-461-1493 | Fax: 713-465-1533

E-Mail: [stoller@stollerusa.com](mailto:stoller@stollerusa.com) | Website: [www.StollerUSA.com](http://www.StollerUSA.com)