

X-TRA POWER PREMIER™



A PLANT GROWTH REGULATOR AND YIELD STIMULANT

F2399
Z-TRAPREMIER

ACTIVE INGREDIENT:

Cytokinin (as kinetin).....0.0075%

INERT INGREDIENTS.....99.9925%

TOTAL.....100.00%

(Contains 0.044 g Cytokinin/pint)

CONTAINS NON-PLANT FOOD INGREDIENT: 0.0075% Cytokinin

GUARANTEED ANALYSIS

Magnesium (Mg).....0.80%

0.80% Chelated Magnesium

Copper (Cu).....0.80%

0.80% Chelated Copper

Manganese (Mn).....0.80%

0.80% Chelated Manganese

Zinc (Zn)3.20%

3.20% Chelated Zinc

(Derived from Magnesium EAHP, Copper EAHP,
Manganese EAHP, Zinc EAHP)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements inside booklet.

Net Weight: 10.35 lbs/gal or 1.24 kg/L

Net Contents:

1 Gal (4 L) 2.5 Gal (10 L)

5 Gal (19 L) 55 Gal (208 L)

Lot Number:

Manufacture Date:

Best if used by:

EPA Reg No. 57538-69

EPA Est No: 57538-TX-2 57538-FL-1 57538-IA-1

FIRST AID

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or are going for treatment. For medical emergencies, call the poison control center at 1-800-222-1222. For general information on this product, call 1-800-920-0131 during the hours of 8-5 EST weekdays, or contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool place and out of direct sunlight.

PESTICIDE DISPOSAL: To avoid wastes, use all of the material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING: For containers 5 gallons or less: Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **For containers 5 gallons or greater: Triple Rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Manufactured by **STOLLER ENTERPRISES, INC.**

9090 Katy Fwy., Suite 400 | Houston, TX 77024 U.S.A.

Toll Free 1-800-539-5283 Tel.: 1(713)461-1493 • Fax: 1(713)465-1533

X-TRA POWER PREMIER™



Stoller

A PLANT GROWTH REGULATOR AND YIELD STIMULANT

ACTIVE INGREDIENT:

Cytokinin(as kinetin).....0.0075%

INERT INGREDIENTS.....99.9925%

TOTAL100.00%

(Contains 0.044 g Cytokinin/pint)

CONTAINS NON-PLANT FOOD

0.0075% Cytokinin

GUARANTEED ANALYSIS

Magnesium (Mg).....0.80%

0.80% Chelated Magnesium

Copper (Cu).....0.80%

0.80% Chelated Copper

Manganese (Mn).....0.80%

0.80% Chelated Manganese

Zinc (Zn).....3.20%

3.20% Chelated Zinc

(Derived from Magnesium EAHP, Copper EAHP, Manganese EAHP, Zinc EAHP)

Information regarding the contents and levels of metals in

this product is available on the internet at

<http://www.aapfc.org/metals.html>

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS INSIDE BOOKLET

NET WEIGHT: 10.35 lbs/gal or 1.24 kg/L

EPA Reg. No. 57538-69

EPA Est. No.: 57538-TX-2, 57538-FL-1, 57538-IA-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin or swallowed. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are any waterproof material. If you want more options, follow instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- waterproof gloves such as polyethylene or polyvinyl chloride material
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

- Users should remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Label updated 02/25/2020

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> · Take off contaminated clothing. · Rinse skin immediately with plenty of water for 15-20 minutes. · Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> · Call a poison control center or doctor immediately for treatment advice. · Have person sip a glass of water if able to swallow. · Do not induce vomiting unless told to do so by the poison control center or doctor. · Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or are going for treatment. For medical emergencies, call the poison control center at 1-800-222-1222. For general information on this product, call 1-800-920-0131 during the hours of 8-5 EST weekdays, or contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu.</p>	
<p>FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.</p>	

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash water or rinsate.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms and in forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours unless wearing the appropriate PPE.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, wear:

- coveralls over long-sleeved shirt and pants,
- waterproof gloves such as polyethylene or polyvinyl chloride material and
- shoes plus socks.

CHEMIGATION[*]

Application and Calibration Techniques for Sprinkler Irrigation

Apply this product only through the following types of irrigation systems: sprinkler including center pivot, traveler, big gun, lateral move, end tow, side (wheel) roll, solid set, or hand move irrigation; furrow; or drip (trickle) irrigation

systems. Do not apply through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Experiment Station specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

[*Not for use in California]

A. Center Pivot, Traveler, Big Gun, Lateral Move, End Tow, and Side (Wheel) Roll Irrigation Equipment:

Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of product for acreage to be covered into same amount of water used during

calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until product has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment:

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of product for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that product will remain in suspension during the injection cycle. Product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until pesticide is cleared from last sprinkler head.

Safety Devices for Sprinkler Chemigation

- (1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check

valve to prevent the flow of fluid back toward the injection pump.

- (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- (5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems Connected to Public Water Sources

- (1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at

least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.

- (2) Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- (3) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- (4) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water

pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Furrow Chemigation[*]

- (1) Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- (2) Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of

fluid back toward the injection pump.

- c. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- e. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- f. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Apply X-TRA POWER PREMIER in sufficient water to penetrate into the root zone without excessive leaching into deeper soil.

[*Not for use in California]

Drip (Trickle) Chemigation[*]

- (1) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- (2) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- (3) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- (5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible

with pesticides and capable of being fitted with a system interlock.

Apply X-TRA POWER PREMIER in sufficient water to penetrate into the root zone without excessive leaching into deeper soil.

[*Not for use in California]

GENERAL USE INSTRUCTIONS

X-TRA POWER PREMIER provides plant essential nutrients Zinc, Manganese, Magnesium, and Copper. X-TRA POWER PREMIER is formulated for efficient uptake by root and leaf tissues of the plant. Mix X-TRA POWER PREMIER with enough water to get thorough coverage of plant surfaces. X-TRA POWER PREMIER is compatible with most fertilizer and pesticide materials, but always conduct a jar test when using an untried combination to ensure compatibility.

CROP USAGE - ALL CROPS LISTED FOR TRANSPLANTING[*] AND SEED BED TREATMENT[*]

Use 32 fl. oz. /acre X-TRA POWER PREMIER per acre (2.4 liters/hectare) or 1 part X-TRA POWER PREMIER to 1000 parts water as a root dip and watering solution when transplanting.

Use 32 fl. oz. /acre X-TRA POWER PREMIER per acre (2.4 liters/hectare) applied to the seedbed at time of seeding or up to 20 days thereafter.

[*Not for use in California]

DILUTION:

For applications by ground application equipment dilute recommended rate of X-Tra Power Premier in a minimum of 10 gallons of water or spray solution, for low volume aerial applications dilute recommended rate in a minimum of 2 gallons of water or spray solution.

MIXING INSTRUCTION:

Follow this mixing order 1. Water 2. X-TRA POWER PREMIER 3. Pesticide. X-TRA POWER PREMIER will disperse in water with little agitation. X-TRA POWER PREMIER is compatible with most fertilizers and pesticides. Always conduct a jar test when using new or untried combinations. The addition of 0.5% (total solution) of nitrogen solution, ammonium sulfate, or low biuret urea may aid leaf absorption.

USE RATES FOR FOLIAR, SOIL, IN-FURROW AND/OR CHEMIGATION[*] APPLICATION:

FOR ALL CROPS LISTED BELOW

Use the higher rate listed in the use rates below by crop, for single planned foliar applications or through in furrow or chemigation (single or multiple) applications. With planned multiple foliar applications, the lower rates in the range below by crop applied multiple times is acceptable.

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
ALFALFA[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	At spring green up with repeat applications within 7 days after cutting each, with repeat sprays at 14 to 21 day intervals.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
APPLES[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: at full pink flower bud. 2nd application: at calyx (petal fall). 3rd application: 3 weeks after 2nd spraying. 4th application: 4 weeks after 3rd spraying.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
ASPARAGUS[*]	16-32 fl. oz./acre (1.2 to 2.4 liters/ hectare)	1st application: spray crowns when growth begins. 2nd application: spray crowns after each cutting.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
BANANAS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	To reduce stress: Apply when stress conditions are anticipated. Rates and timing must be determined for each site. Make applications at least 14 days apart using ground sprayers, aerial sprayers, or by plant injection.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
BARLEY[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 14-21 days after planting. 2nd application: Late tiller stage. Repeat applications may be made every 14-21 days through heading.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
ALL BEAN CROPS[*] (Green Beans[*], Snap Beans[*], Dry Edible Beans[*], Lentils[*])	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	May apply in furrow at planting. Foliar Applications: 1st application: 4-5 inch stage. 2nd application: at early bloom. 3rd application: at early pod set.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
BEETS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: at tuber initiation. 2nd application: 2-3 weeks after 1st spraying.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
CANOLA[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	May apply in furrow at planting. Application: Apply 14-21 days after planting, with repeat application every 7-14 days	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
CARROTS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: at tuber initiation. 2nd application: 14-21 days after first spraying.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
CELERY[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	Application: Use 16 fl. oz. /acre (1.2 liters/hectare) within 7 days after transplanting, repeat applications may be made 10 to 21 day intervals	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
CITRUS CROPS[*] (Oranges[*], Grapefruit[*], Lemons[*], Limes[*], Tangerines[*])	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: pre-bloom. 2nd application: at calyx (petal fall). 3rd application: 3 weeks after 2nd spraying. 4th application: 4 weeks after 3rd spraying.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
CORN[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	May apply in furrow at planting. Foliar applications: 1st application: At V3-V5 growth stage. 2nd application: 10-14 days pre tassel or post tassel at R2 plant development stage.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
COTTON[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	May apply in furrow at planting. Foliar applications: 1st application: 30-40 days after planting with repeat applications at 14 to 21 day intervals.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
CRUCIFEROUS CROPS[*] (Cabbage, Broccoli, Cauliflower, Brussels Sprout)	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 3 to 4-inch stage. Repeat at 10 to 14 day intervals.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
CUCURBITS[*] (Cucumbers[*], Muskmelon[*], Cantaloupe[*], Watermelon[*], Honey Dew[*], Squash[*], etc.)	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 20-30 days after planting. 2nd application: at early bloom. 3rd application: start of fruit development.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
GRAPES[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: between leaf-out and pre-bloom. 2nd application: at calyx (petal fall). 3rd application: 30 days before harvest.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
OATS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: Late tiller growth stage 2nd application: Late boot stage to flag leaf stage.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
OLIVE[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	Every 7 to 21 days from bud break through harvest.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
ONIONS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 7-14 days after transplanting, with repeat applications every 14-21 days up to harvest.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
PEACHES[*] AND NECTARINES[*]	16-32 fl. oz./acre (1.2 to 2.4 liters/hectare)	1st application: at pre-bloom. 2nd application: at calyx (petal fall). 3rd application: 3 weeks after 2nd spraying. 4th application: 4 weeks after 3rd spraying.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
PEANUTS[*]	16-32 fl. oz./acre (1.2 to 2.4 liters/hectare)	May apply in furrow at planting. Foliar Applications: 1st application: at pegging, with repeat applications made every 10-14 day intervals continuing up to 20-30 days pre-harvest.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
PEAS[*]	16-32 fl. oz./acre (1.2 to 2.4 liters/hectare)	May apply in furrow at planting. Foliar applications: 1st application: 3 to 4-inch stage. 2nd application: Pre-bloom. 3rd application: at early pod set.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
PEPPERS[*] AND EGGPLANT[*]	16-32 fl. oz./acre (1.2 to 2.4 liters/hectare)	1st application: just prior to 1st bloom. 2nd application: 10 days after 1st spraying. 3rd application: 10 days after 2nd spraying.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
PINEAPPLE[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	To reduce plant stress [^] : Apply to vegetative growth according to climate and crop needs at the site of proposed application. To improve fruit growth [^] : Apply post bloom according to climate and crop needs at the site of proposed application. [^] Allow at least 14 days between applications.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
POMEGRANATE[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	Every 7 to 21 days from bud break through harvest.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
POTATOES[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: at tuber set (5-6 weeks after planting). 2nd application: at full blossom (14-21 days after 1st application)	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
RICE[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: at 2 to 5 leaf stage with repeat application 14 to 21 days after.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year
RYE[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 14-21 days after planting or for fall planted crops apply at spring green up. 2nd application: Late tiller stage. With repeat applications made every 14-21 days through heading.	64 fl. oz./acre (4.8 liters/ hectare) per application; 128 fl. oz./acre (9.6 liters/ hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
SORGHUM (MILO) [*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/hectare)	May apply in furrow at planting. 1st application: At V3-V5 growth stage. 2nd application: 10-14 days pre heading	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
SOYBEANS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/hectare)	May apply in furrow at planting. Foliar Application: Apply beginning at V3 growth stage, with repeat application every 14-21 days up to R5	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
SPINACH[*] AND LETTUCE[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/hectare)	Application: 21-28 days after planting or 7-14 days after transplanting.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
STRAWBERRIES[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/hectare)	Application: Starting 7-14 days after planting or pre-bloom, with repeat applications made every 10-14 days.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
SUGAR BEETS[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/hectare)	1st application: at tuber initiation. 2nd application: 2-3 weeks after 1st spraying.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
SUNFLOWER[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/hectare)	May apply in furrow at planting. 1st application: 3-4 leaf growth stage. 2nd application: at bud stage. 3rd application: at flowering	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year

(continued)

CROP	USE RATE	APPLICATION	MAXIMUM APPLICATION RATES
SWEET POTATOES[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: at tuber set (5-6 weeks after planting). 2nd application: at full blossom (14-21 days after 1st application)	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
TOMATOES[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 7-14 days after transplanting. 2nd application: 14-21 days after 1st spray. 3rd application: 14-21 days after 1st bloom.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
TREE NUT CROPS[*] (Almond[*], Pecan[*], Pistachio[*], Walnut[*])	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: pre-bloom. 2nd application: at calyx (petal fall). 3rd application: 14-21 days after 2nd spraying. 4th application: 21-28 days after 3rd spraying.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
WHEAT[*]	16-32 fl. oz. /acre (1.2 to 2.4 liters/ hectare)	1st application: 14-21 days after planting or for fall planted crops apply at spring green up. 2nd application: Late tiller stage. With repeat applications made every 14-21 days through heading.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year
ORNAMENTAL TREES AND HERBACEOUS PLANTS[*]		Apply 32 fl. oz. per acre as a foliar spray when growth begins in the early spring. Apply 32 fl. oz. per acre at the end of summer to maintain color through autumn and aid in winter survival.	64 fl. oz./acre (4.8 liters/hectare) per application; 128 fl. oz./acre (9.6 liters/hectare) per year

[*Not for use in California]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool place and out of direct sunlight.

PESTICIDE DISPOSAL: To avoid wastes, use all of the material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING: **For containers 5 gallons or less: Triple Rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **For containers 5 gallons or greater: Triple Rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

WARRANTY

To the fullest extent permitted by law, neither the manufacturers nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.

Manufactured by:

STOLLER ENTERPRISES, INC.

9090 Katy Frwy., Suite 400
Houston, TX 77024 U.S.A.
Toll Free 1-800-539-5283
Tel.: 1(713) 461-1493
Fax: 1(713) 465-1533

X-TRA POWER PREMIER™



A PLANT GROWTH REGULATOR AND YIELD STIMULANT

ACTIVE INGREDIENT:

Cytokinin(as kinetin).....0.0075%

INERT INGREDIENTS.....99.9925%

TOTAL100.00%

(Contains 0.044 g Cytokinin/pint)

CONTAINS NON-PLANT FOOD

0.0075% Cytokinin

GUARANTEED ANALYSIS

Magnesium (Mg).....0.80%

0.80% Chelated Magnesium

Copper (Cu).....0.80%

0.80% Chelated Copper

Manganese (Mn).....0.80%

0.80% Chelated Manganese

Zinc (Zn).....3.20%

3.20% Chelated Zinc

(Derived from Magnesium EAHP, Copper EAHP, Manganese EAHP, Zinc EAHP)

Information regarding the contents and levels of metals in

this product is available on the internet at

<http://www.aapfc.org/metals.html>

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

SEE ADDITIONAL PRECAUTIONARY STATEMENTS INSIDE BOOKLET

NET WEIGHT: 10.35 lbs/gal or 1.24 kg/L

EPA Reg. No. 57538-69

EPA Est. No.: 57538-TX-2, 57538-FL-1, 57538-IA-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin or swallowed. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are any waterproof material. If you want more options, follow instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- waterproof gloves such as polyethylene or polyvinyl chloride material
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

- Users should remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Label updated 02/25/2020