

# Stoller®

# Root Feed Dry

## 9-0-5

### GUARANTEED ANALYSIS

Total Nitrogen (N).....	9.0%
9% Urea Nitrogen	
Soluble Potash (K <sub>2</sub> O).....	5.0%
Calcium (Ca).....	12.9%
Magnesium (Mg).....	1.2%
Boron (B).....	0.12%
Molybdenum (Mo).....	0.03%

Derived from urea, calcium chloride, potassium chloride, magnesium chloride, boric acid and sodium molybdate.

**STOP, READ THE LABEL BEFORE  
USING THE PRODUCT  
CAUTIONS AND WARNINGS  
KEEP OUT OF REACH OF CHILDREN**

This product must be used based on the instructions stated on this label, use a mask and rubber boots when applying this product. If inhaled remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention. If in eyes wash with plenty of water. Call physician if irritation persists. If on skin wash with plenty of soap and water for at least 15 minutes. Get medical attention if irritation persists. If ingested do not induce vomiting but call a physician and take patient to an emergency facility. May be harmful if absorbed through skin. May cause eye irritation. Avoid contact with skin, eyes or clothing. Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves and shoes plus socks. Follow manufacturer instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENVIRONMENTAL HAZARDS:

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

### STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Store in a cool place and out of direct sunlight. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NET CONTENTS:

### DIRECTIONS FOR USE

#### Vegetables

DOSES (Kg/Ha): 25 – 40

Period of Application: Apply as stated every 7 to 14 days after the Annual transplant or from the green, in case of a direct planting through irrigation water. Continue the applications with intervals of 7 to 10 days, up to 30 to 15 days before the harvest. Begin after seedling or transplanting when plants are well established and continue until harvest is completed.

#### Fruit Trees

DOSES (Kg/Ha): 0 – 40

Period of Application: Begin the applications when sprouting begins before green tissue appears. Also, begin 3 weeks before flower buds appear and continue applications until 3 weeks after harvest or desired maturity. Continue intervals from 10 -14 days up to 4 or 8 weeks before the harvest.

### MIXING INSTRUCTIONS:

Do not mix with fertilizers containing phosphorus or sulfur. Many pesticides may be mixed and applied with any solution containing Stoller Root Feed Dry. Mixing sequence: 1. Water 2. Stoller Root Feed Dry 3. Pesticide or other fertilizer. Always conduct a "jar test" utilizing all products to be mixed in proper proportion in order to establish compatibility.

ORDER OF MIXTURE: 1. Water 2. Stoller Root Feed Dry 3. Chemical Product

### CAUTION:

This product contains BORON and its use on any crop other than those recommended on this label may result in serious injury to the crop.

### CAUTION:

This product contains MOLYBDENUM. The application of fertilizers with Molybdenum may result in forage crops containing levels of Molybdenum, which are toxic to ruminant animals. Keep out of reach of children.

### CAUTION:

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>.

Z-SRFDRY

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 | Houston, TX 77024 | Phone: 1-800-539-5283 or 713-461-1493

