

# Harvest More<sup>®</sup> 20-20-20



## SPECIMEN

### 20-20-20 GUARANTEED ANALYSIS

Total Nitrogen (N).....	20%
4% Ammoniacal Nitrogen	
16% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....	20%
Soluble Potash (K <sub>2</sub> O).....	20%
Boron (B).....	0.02%
Cobalt (Co).....	0.002%
0.002% Chelated Cobalt	
Copper (Cu).....	0.05%
0.05% Chelated Copper	
Manganese.....	0.05%
0.05% Chelated Manganese	
Molybdenum (Mo).....	0.005%
Zinc (Zn).....	0.1%
0.1% Chelated Zinc	

Plant nutrients derived from urea, monoammonium phosphate, potassium chloride, boric acid, sodium molybdate, cobalt EDTA, copper EDTA, manganese EDTA and zinc EDTA.

F2399

Z-HM202020  
Rev: 22F27

### NET WEIGHT

☐ 55 lb (25 kg)

### CAUTION

KEEP OUT OF REACH OF CHILDREN, MAY BE HARMFUL IF SWALLOWED, INHALED OR IN CONTACT WITH SKIN; MAY CAUSE EYE IRRITATION. MAY CAUSE DAMAGE THROUGH PROLONGED EXPOSURE.

#### PRECAUTIONS:

Do not breathe dust/fume/gas/mist/vapors/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in well-ventilated area. Avoid release to environment. Wear protective gloves/protective clothing/eye protection/face protection. Keep container tightly closed. Keep cool and protected from sunlight. If medical advice is needed, have product container or label at hand. For more details regarding the safe use/handling, storage and disposal of this product as well as additional hazards and first aid measures, please obtain the current Safety Data Sheet at [www.stollerusa.com](http://www.stollerusa.com).

### STORAGE AND DISPOSAL

Store bags tightly closed in a cool/well-ventilated dry place protected from sunlight and humidity. Store locked up. Do not contaminate water, food or feed by storage or disposal. Do not apply directly to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

**WARNING:** This product contains boron at a level that may be detrimental to growing plants. Use of boron on crops other than those recommended may result in serious injury to those crops.

**WARNING:** The application of fertilizing materials containing molybdenum may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>.

*Always read and follow label instructions.*

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | [www.stollerusa.com](http://www.stollerusa.com) | 1-800-539-5283 or 713-461-1493 | ©2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.

---

## GENERAL INFORMATION

---

HARVEST MORE® 20-20-20 is a foliar applied micronutrient and is non-phytotoxic when used as directed. It is an acidification agent for lowering high pH in spray tank water to decrease the hydrolysis of most pesticides and is an effective dispersing agent when applied with other foliar spray chemicals.

---

## DIRECTIONS FOR USE

---

It may be a violation of law to use this product inconsistent with its labeling.

---

## RECOMMENDED USES

---

**FOR TRANSPLANT APPLICATION:** Use 5 pounds per acre with a dissolution requirement of 100 gallons of water (rate of use per acre should be 5% of total tank capacity. With a 50-gallon spray tank, use 2.5 pounds per acre).

**FOR FOLIAR APPLICATION:**

**Field Crop:** Alfalfa, barley, canola, corn, cotton, flax, milo, oats, soybeans, sugar beets, sugar cane and wheat. Apply 1 to 2 pounds per 100 gallons of spray solution. Do not apply more than 2.5 to 5 pounds per acre per application. Apply as needed throughout the season.

**Vegetables:** All beans, broccoli, cabbage, cauliflower, carrots, celery, lettuce, melons, onions, peppers, potatoes, peanuts, sweet corn, spinach, strawberries, tomatoes, turnips and watermelons. Apply 10 pounds per 100 gallons of spray solution. Apply every 7 to 10 days or as needed starting pre-flowering to end of harvest. Full coverage sprays work best.

**Tree and Vine Crops:** Almonds, apples, apricots, avocados, cherries, citrus, filberts, grapes, nectarines, pears, peaches, pecans, plums, prunes, and walnuts. Apply 10 pounds in 100 gallons of water or spray solution. Apply pre-bloom and post-bloom as needed avoid applications during bloom. Full coverage sprays work best.

**House Plants-Nursery and Greenhouses:** For gardenias, carnations, African violets, etc., mix 1 tablespoon in 1 gallon of water. Apply as a drench to soil or spray twice monthly. For African violets apply as a drench only. Use 10 pounds per 100 gallons on plants, shrubs, flowers and trees for direct spray to foliage. For transplant slurry, add ¼ lb or ½ standard cupful per 3-gallon bucket of puddled soil. Use 1 ¾ pounds in a sprinkler system per 100 gallons water every 2 to 3 weeks.

**Lawns, Golf Courses, and Turf Areas:** For Kentucky and Merion blue, bent, Bermuda and Zoysia grasses. The type of grass color and rate of growth will depend on the frequency of application. Harvest More can be combined with fungicides and/or soil insect treatments. Apply 1 pound per 100 gallons of water per 1000 sq. ft. Apply monthly, starting February or March. Frequent applications may be necessary for desirable color and growth, especially on Bermuda and Zoysia Grasses. Apply with watering can or sprayer.

NOTE: Grasses growing in shade require approximately one-half as much plant food as when grown in full sun.

**FOR GENERAL APPLICATION:** Use 10 pounds per 100 gallons water or spray solution. For most crops apply every 7 to 10 days.

**MIXING INSTRUCTIONS:** Follow this mixing order – (1) Water (2) HARVEST MORE® 20-20-20 (3) Pesticide. HARVEST MORE® 20-20-20 will disperse in water with little agitation. HARVEST MORE® 20-20-20 is compatible with many fertilizers and pesticides. Always conduct a jar test when using new or untried combinations. The addition of 0.5% (total solution) of nitrogen solution, ammonium sulfate, or low biuret urea may aid leaf absorption.

NOTICE: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller make any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

*Always read and follow label instructions.*

**Manufactured and Guaranteed by Stoller Enterprises, Inc.**

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | www.stollerusa.com | 1-800-539-5283 or 713-461-1493 | ©2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.